

Label-free identification of single cell-derived vesicles by Raman microspectroscopy

Edwin van der Pol^{1,2}


Chi Hau², Cees Otto³, Aufried Lenferink³,
Auguste Sturk², Ton van Leeuwen¹,
and Rienk Nieuwland²

February 6th, 2013




¹Biomedical Engineering and Physics; ²Laboratory Experimental Clinical Chemistry, Academic Medical Center, Amsterdam, The Netherlands; ³Medical Cell BioPhysics, University of Twente, Enschede, the Netherlands

Introduction – motive





Introduction – cell-derived vesicles



- (tumor) cells release vesicles:
 - spherical particles with phospholipid bilayer
- specialized functions

Introduction – cell-derived vesicles



- detection and isolation cumbersome
 - small size (< 500 nm)
 - low refractive index (~1.4)
 - fluorescent antibody labeling involves practical problems


Goal



erythrocyte
vesicles




platelet
vesicles



Human pancreatic
adenocarcinoma
(BxPC-3) vesicles

- distinguish tumor (BxPC-3) vesicles from erythrocyte and platelet vesicles directly in solution *without labeling* by Raman microspectroscopy

Method 1 – Single particle tracking



$$\langle x^2 \rangle = \frac{2k_B T \tau}{3\pi\eta d}$$

x : displacement

k_B : Boltzmann constant

T : temperature

t : observation time

η : viscosity



d : diameter

Method 1 – Single particle tracking




- single particle tracking applied to determine size and concentration of vesicles


Results – Single particle tracking




Method 2 - Raman microspectroscopy



Methods - Raman microspectroscopy




PBS




Tumor (BxPC-3) vesicle

Methods - Raman microspectroscopy





Erythrocyte vesicles





Erythrocyte vesicles



Methods - Raman spectrum of PBS





Methods - Raman spectrum of *single* tumor (BxPC-3) vesicle





Methods - Raman spectrum of *single* tumor (BxPC-3) vesicle





Methods - Raman spectrum of *single* tumor (BxPC-3) vesicle






Results - Raman spectrum of *single erythrocyte vesicles*





Results - Raman spectrum of *single* platelet vesicles



Results - Raman spectrum of *single* BxPC-3 vesicles



Results - summary



Conclusion and discussion

- distinguished single tumor (BxPC-3) vesicles from erythrocyte and platelet vesicles directly in solution *without labeling*
- improvements
 - simultaneous sizing by single particle tracking
 - optimize isolation
- develop high throughput system
 - microfluidics
 - multiple probes
 - Coherent anti-Stokes Raman spectroscopy

Acknowledgements

- Anita Böing
- Anita Grootemaat
- Aufried Lenferink
- Cees Otto
- Chi Hau
- Guus Sturk
- Henk van Veen
- Rienk Nieuwland
- Ton van Leeuwen

